

Patient Alert Card
ZENTIVA

Xanirva 20 mg (tick the prescribed dose)

Keep this card with you at all times

Present this card to every physician or dentist prior to treatment

I am under anticoagulation treatment with Xanirva (rivaroxaban)

Name:

Address:

Birth date:

Weight:

Other medicines / conditions:

In case of emergency, please notify:

Doctor's name:

Doctor's phone:

Doctor's stamp:

Please also notify:

Name:

Phone:

Relationship:

Information for health care providers:

- INR values should not be used as they are not a dependable measure of the anticoagulant activity of **Xanirva**.

What should I know about Xanirva?

- Xanirva thins the blood, which prevents you from getting dangerous blood clots.
- Xanirva must be taken exactly as prescribed by your doctor. To ensure optimal protection from blood clots, **never skip a dose**.
- You must not stop taking Xanirva without first talking to your doctor as your risk of blood clots may increase.
- Tell your health care provider about any other medicines you are currently taking, took recently or intend to start taking, before you start Xanirva.
- Tell your health care provider that you are taking Xanirva before any surgery or invasive procedure.

When should I seek advice from my health care provider?

When taking a blood thinner such as Xanirva it is important to be aware of its possible side effects. Bleeding is the most common side effect. Do not start taking Xanirva if you know you are at risk of bleeding, without first discussing this with your doctor. Tell your health care provider straight away if you have any signs or symptoms of bleeding such as the following:

- pain
- swelling or discomfort
- headache, dizziness or weakness
- unusual bruising, nosebleeds, bleeding of gums, cuts that take a long time to stop bleeding
- menstrual flow or vaginal bleeding that is heavier than normal
- blood in your urine which may be pink or brown, red or black stools
- coughing up blood, or vomiting blood or material that looks like coffee grounds

How do I take Xanirva?

- To ensure optimal protection, Xanirva
 - 20 mg must be taken with food

Sources of Global document:

- SmPC – for film-coated tablets (CZ/H/0834-0835) and hard capsules (CZ/H/0836-0837) approved 29-Apr-2022.
- RMP – version 2.0 for film-coated tablets (CZ/H/0834-0835, approved 28-Aug-2022) and version 1.2 for hard capsules (CZ/H/0836-0837, approved 22-Mar-2022)
- Educational material of the reference product Xanirva:
 - Patient Alert Card (Jan-2019)